

REFERENCES

Training Tools – Explaining Contraception and the Birth Control Method Options chart

May 26, 2016

Reference for effectiveness rates: Trussell J. Contraceptive failure in the United States. *Contraception* 2011; 83: 397-404.

Long Acting Reversible Contraception (LARC): IUDs and implant

Diedrich JT, Desai S, Zhao Q, Secura G, Madden T, Peipert JF. Association of short-term bleeding and cramping patterns with long-acting reversible contraceptive method satisfaction. *Am J Obstet Gynecol*. 2015 (212) 1: 50.e1-50 e8.

Dhamangaonkar PC, Anuradha K, Saxena A. Levonorgestrel intrauterine system (Mirena): An emerging tool for conservative treatment of abnormal uterine bleeding. *J Midlife Health*. 2015 (6) 1:26.

Gemzell-Danielsson K, Inki P, Boubli L, et al. Bleeding pattern and safety of consecutive use of the levonorgestrel-releasing intrauterine system (LNG-IUS)--a multicentre prospective study. *Hum Reprod*. 2010;25(2):354-9.

Heikinheimo O, Schmelter T, Gemzell-Danielsson K. Bleeding pattern and user satisfaction in second consecutive levonorgestrel-releasing intrauterine system users: results of a prospective 5-year study. *Hum Reprod*. 2014; 29 (6): 1182-1188.

Hidalgo M, Bahamondes L, Perrotti M, et al. Bleeding patterns and clinical performance of the levonorgestrel-releasing intrauterine system (Mirena) up to two years. *Contraception*. 2002;65(2):129-32.

Larsson G, Milsom I, Jonasson K, et al. The long-term effects of copper surface area on menstrual blood loss and iron status in women fitted with an IUD. *Contraception*. 1993; 48(5):471-80.

Mansour D, Korver T, Marintcheva-Petrova M, Fraser IS. The effects of Implanon on menstrual bleeding patterns. *Eur J Contracept Reprod Health Care*. 2008; (Suppl 1): 13-28.

Milsom I, Andersson K, Jonasson K, et al. The influence of the Gyne-T 380S IUD on menstrual blood loss and iron status. *Contraception*. 1995; 52:175–179.

Milsom I, Rybo G, Lindstedt G. The influence of copper surface area on menstrual blood loss and iron status in women fitted with an IUD. *Contraception*. 1990; 41(3):271-81.

Rowe PP, Farley T, Peregoudov A, et.al. Safety and efficacy in parous women of a 52-mg levonorgestrel-medicated intrauterine device: a 7-year randomized comparative study with the TCu380A. 2016: doi: 10.1016/j.contraception.2016.02.024

Weisberg E, Bateson D, McGeechan K, Mohapatra L. A three-year comparative study of continuation rates, bleeding patterns and satisfaction in Australian women using a subdermal contraceptive implant or progestogen releasing-intrauterine system. *Eur J Contracept Reprod Health Care*. 2013; 19 (1): 5-14.

REFERENCES

Training Tools – Explaining Contraception and the Birth Control Method Options chart

May 26, 2016

Hormonal Methods: Oral Contraception, Vaginal Ring, Patch, Injection

Arias RD, Jain JK, Brucker C, Ross D, Ray A. Changes in bleeding patterns with depot medroxyprogesterone acetate subcutaneous injection 104 mg. *Contraception*. 2006; 45: 234-238

Audet M-C, Moreau M, Koltun WD, et al. Evaluation of contraceptive efficacy and cycle control of a transdermal contraceptive patch vs an oral contraceptive. A randomized controlled trial. *JAMA*. 2001; 285:2347-54.

Burkman R, Schlesselman JJ, Ziemann M. Safety concerns and health benefits associated with oral contraception. *Am J Obstet Gynecol*. 2004; 190(Suppl):S5-S22.

Bjarnadottir RI, Tuppurainen M, Killick SR. Comparison of cycle control with a combined contraceptive vaginal ring and oral levonorgestrel/ethinyl estradiol. *Am J Obstet Gynecol*. 2002; 186(3):389-95.

Brunei V, Pontello V, Luisi S, et al. An open-label, multicentre trial to evaluate the vaginal bleeding pattern of the combined contraceptive vaginal ring NuvaRing. *Eur J Obstet Gynecol Reprod Biol*. 2008;39(1):65-71.

Davis A, Godwin A, Lippman J, et al. Triphasic norgestimate ethinyl estradiol for treating dysfunctional dysphoric disorder. *Obstet Gynecol*. 2005; 106:492-501

Dieben TO, Roumen FJ, Apter D. Efficacy, cycle control, and user acceptability of a novel combined contraceptive vaginal ring. *Obstet Gynecol*. 2002; 100(3):585-93.

Dragoman MV, Gaffield ME. The safety of subcutaneously administered depot medroxyprogesterone acetate (104mg/0.65mL): A systematic review. *Contraception*. In press

Gallo MF, Lopez LM, Grimes DA, Schulz KF, Helmerhorst FM. Combination contraceptives: Effects on weight. *Cochrane DB Syst Rev*. 2006 (1):CD003987.

Jacobstein R, Polis C. progestin-only contraception: Injectables and implants. *Best Pract Res Cl Ob*. 2014; 28 (6): 795-806.

Kwiecien M, Edelman A, Nichols MD, et al. Bleeding patterns and patient acceptability of standard or continuous dosing regimens of a low-dose oral contraceptive: a randomized trial. *Contraception*. 2003; 67:9-13.

Larsson G, Milsom I, Lindstedt G, et al. The influence of a low-dose combined oral contraceptive on menstrual blood loss and iron status. *Contraception*. 1992; 46 (4):327-34

Lopez LM, Grimes DA, Gallo MF, Stockton LL, Schulz KF. Skin patch and vaginal ring versus combined oral contraceptives for contraception. *Cochrane DB Syst Rev*. 2013; 4: CD003552.

Le YL, Rahman M, Berenson A. Early weight gain predicting later weight gain among depot medroxyprogesterone acetate users. *Obstet Gynecol*. 2009; 114 (2, Part 1):279-84.

REFERENCES

Training Tools – Explaining Contraception and the Birth Control Method Options chart

May 26, 2016

McCann MF, Potter LS. Progestin-only contraception: a comprehensive review. *Contraception*. 1994; 50:S1-195.

Nelson AL. Counseling issues and management of side effects for women using depot medroxyprogesterone contraception. *J Reprod Med*. 1996; 41 (Suppl 5):391.

Redmond GP, Olson WH, Lippman JS, et al. Norgestimate and ethinyl estradiol in the treatment of acne vulgaris: a randomized, placebo-controlled trial. *Obstet Gynecol*. 1997; 89:615-22

Risser WL, Geftter LR, Barratt MS, et al. Weight change in adolescents who used hormonal contraception. *J Adolesc Health*. 1999;24(6):433-6.

Sibai BM, Odlind V, Meador ML, Shangold GA, Fisher AC, Creasy GW. A comparative and pooled analysis of the safety and tolerability of the contraceptive patch (Ortho Evra/Evra). *Fertil Steril*. 2002; 77 (2 Suppl 2) S19-26.

Veisi F, Zangeneh M. Comparison of two different injectable contraceptive methods: Depo-medroxy progesterone acetate (DMPA) and cyclofem. *J Family Reprod Health*. 2013; (3): 109-113.

Zieman M, Guillebaud J, Weisberg E, et al. Contraceptive efficacy and cycle control with the Ortho Evra/Evra transdermal system: the analysis of pooled data. *Fertil Steril*. 2002; 77(2 Suppl 2):S13-S18.

OTHER AVAILABLE METHODS

Abstinence

World Health Organization. Defining sexual health: report of a technical consultation on sexual health, 28-31 January 2002, Geneva: WHP, 2006.

ACOG Committee Opinion No. 417: Addressing health risks of noncoital sexual activity. *Obstet Gynecol*. 2008; 112:735-737.

Kirby D. Emerging Answers 2007: research findings on programs to reduce teen pregnancy and sexually transmitted disease. Washington, DC: The National Campaign to Prevent Teen and Unplanned Pregnancy, 2007.

Breastfeeding – [Lactational Amenorrhea (LAM)]

Institute for Reproductive Health - http://irh.org/projects/fam_project/lactational-amenorrhea-method-lam/

Lactational Amenorrhea Method: A Reference Guide for Service Providers (2009) -http://irh.org/wp-content/uploads/2013/04/LAM_ReferenceGuide_JAN10.pdf

REFERENCES

Training Tools – Explaining Contraception and the Birth Control Method Options chart

May 26, 2016

AE Peterson, R Pérez-Escamilla, MH Labboka, V Hight, Multicenter study of the lactational amenorrhea method (LAM) III: effectiveness, duration, and satisfaction with reduced client–provider contact, *Contraception*, November 2000 Volume 62, Issue 5, Pages 221–230

Arévalo, M., Jennings V., and Sinai I. (2003). “Application of simple fertility awareness–based methods of family planning to breastfeeding women”. *Fertility and Sterility* 80(5): 1241-1248.

Kazi A, Kennedy KI, Visness CHM and Khan T. 1995. Effectiveness of the Lactational Amenorrhea Method in Pakistan. *Fertil Steril* 64: 717–723.

Kennedy K, Labbok MH and Van Look PFA. 1996. Consensus statement: Lactational amenorrhea method for family planning. *Int J Gynecol Obstet* 54: 55–57.

Labbok MH, Hight-Laukaran V, Peterson AE, Fletcher V, von Hertzen H, Van Look PF. Multicenter study of the Lactational Amenorrhea Method (LAM): I. Efficacy, duration, and implications for clinical application. *Contraception*. 1997 Jun;55 (6):327-36.

Diaphragm

Trussell J, Strickler J, Vaughan B. Contraceptive efficacy of the diaphragm, the sponge and the cervical cap. *Fam Plann Perspect* 1994; 26(2):66-72.

Emergency Contraceptive Pills (ECP)

The Emergency Contraception Website

<http://ec.princeton.edu/emergency-contraception.html>

Trussell, J, Raymond, EG, Cleland, K. *Emergency Contraception: A Last Chance to Prevent Unintended Pregnancy*. March 2016

<http://ec.princeton.edu/questions/ec-review.pdf>

Female Condom

Trussell J, Sturgen K, Strickley J, Dominik R. Comparative contraceptive efficacy of the female condom and other barrier methods. *Fam Plann Perspect*. 1994; 26:66-72.

Female Sterilization

U.S. Selected Practice Recommendations for Contraceptive Use, 2013

<http://www.cdc.gov/reproductivehealth/unintendedpregnancy/usspr.htm>

Fertility Awareness-Based (FAB) Methods

Fertility Awareness-Based Methods (General)

P. Frank-Herrmann, J. Heil, C. Gnoth, E. Toledo, S. Baur, C. Pyper, E. Jenetzky, T. Strowitzki and G. Freundl, The effectiveness of a fertility awareness based method to avoid pregnancy in relation to a

REFERENCES

Training Tools – Explaining Contraception and the Birth Control Method Options chart

May 26, 2016

couple's sexual behavior during the fertile time: a prospective longitudinal study. **Human Reproduction**. 2007. Volume 22, Issue 5 Pp. 1310-1319.

Sinai, I., and M. Arevalo. It's all in the timing: Coital frequency and fertility awareness-based methods of family planning. *Journal of Biosocial Science*. 38 (2005): 763-777.

Gribble JN, Lundgren RI, Velasquez C, Anastasi EE. Being strategic about contraceptive introduction: the experience of the Standard Days Method®. *Contraception*. 2008;77 (3): 147–154.

Fehring, R., Schneider, M., Raviele, K, Rodriguez, D., & Pruszynski, J. (2013). Randomized comparison of two Internet-supported fertility awareness based methods of family planning, *Contraception*, 88(1): 24-30.

Trussell J, Grummer-Strawn L. Contraceptive failure of the ovulation method of periodic abstinence. *Fam Plann Perspect*. 1990 Mar-Apr;22(2):65-75.

Trussell, J. and K. Kost. 1987. Contraceptive failure in the United States: A critical review of the literature. *Studies in Family Planning*. (18) 237-283.

World Health Organization. 1981. A prospective multicenter trial of the ovulation method of natural family planning. II The Effectiveness Phase. *Fertility and Sterility*. (36) 152.

Standard Days Method

Sinai I, Lundgren RI, Gribble JN. Continued use of the Standard Days Method(R). *J Fam Plann Reprod Health Care*. 2012;38 (3): 150–156.

Arevalo, M., I. Sinai, and V. Jennings. 1999. A fixed formula to define the fertile window of the menstrual cycle as the basis of a simple method of natural family planning. *Contraception*. (60) 357-360.

Arevalo, M. V. Jennings, and I. Sinai. 2002. Efficacy of a new method of family planning: the Standard Days Method. *Contraception*. (65) 333-338.

Sinai, I., V. Jennings, and M. Arevalo. 2004. The importance of screening and monitoring: the Standard Days Method and cycle regularity. *Contraception*. (69) 201-206.

TwoDay Method

Sinai, I., Jennings, V., Arévalo, M. (1999). The TwoDay Algorithm: a new algorithm to identify the fertile time of the menstrual cycle. *Contraception*. 60(2):65-70.

Arévalo, M., Jennings V., Nikula M., and Sinai I. (2004). Efficacy of the new TwoDay Method of family planning. *Fertility and Sterility* 82(4):885-892.

Jennings, V., Sinai, I., Sacieta, L., Lundgren, R. (2011). TwoDay Method: a Quick Start Approach. *Contraception*. 84(2):144-9.

REFERENCES

Training Tools – Explaining Contraception and the Birth Control Method Options chart

May 26, 2016

Fertility Awareness Information

Fehring R., & Schneider M. (2008). Variability of the fertile phase of the menstrual cycle. *Fertility and Sterility*, 90(4), 1232-1235.

Wilcox, A. J., Baird, D. D., Dunson, D. B., McConaughey, D. R., Kesner, J. S., & Weinberg, C. R. 2004. On the frequency of intercourse around ovulation: evidence for biological influences. *Human Reproduction*. 19(7), 1539-1543.

Wilcox, A.G., C.R. Weinberg, and D. Baird. 1998. Post-ovulatory aging of the human oocyte and embryo failure. *Human Reproduction*. (13) 394-397.

Odeblad, E. (1997), Cervical Mucus and their Functions. *Journal of the Irish College of Physicians and Surgeons* 26:27-32.

Ecochard, R. 2005. Heterogeneity: The masked part of reproductive technology success rates. *Rev. Epidemiologique de Sante Publique*. (53) 2S 107-117.

Male Condom

Centers for Disease Control and Prevention *Condom Fact Sheet In Brief*
<http://www.cdc.gov/condomeffectiveness/docs/condomfactsheetinbrief.pdf>

Male Sterilization

U.S. Selected Practice Recommendations for Contraceptive Use, 2013
<http://www.cdc.gov/reproductivehealth/unintendedpregnancy/usspr.htm>

Spermicides and Sponge

Harwood B, Meyn LA, Ballagh SA, Raymond EG, Archer DF, Creinin MD. Cervico-vaginal colposcopic lesions associated with 5 nonoxynol-9 vaginal spermicide formulations. *Am J Obstet Gynecol*. 2008; 198:32 e31-37.

Withdrawal

Jones RK, Lindberg LD, Higgins JA. Pull and pray or extra protection? Contraceptive strategies involving withdrawal among US adult women. *Contraception*. 2014; 90:416-421.

Jones RK, Fennell J, Higgins JA, Blanchard K. Better than nothing or savvy risk-reduction practice? The importance of withdrawal. *Contraception*. 2009; 79:407-410.